PROVIDEAN BH RAB LIQUIDA

RABIES VACCINE

Liquid

REG. SAN: 4A-11083-AGROCALIDAD

Formula:

Each 1 mL dose of vaccine contains:
Inactivated Rabies Virus ≥ 1 IU –
N.I.H./dose for dogs and cats.

INDICATIONS:

Rabies Vaccine for Dogs and Cats

ROUTE OF ADMINISTRATION: Subcutaneous

SPECIES AND DOSAGE:



DOSAGE
Administer 1 mL per animal subcutaneously.

SUGGESTED VACCINATION SCHEDULE :

For animals from 12 weeks of age. Administer a booster dose annually. The vaccination schedule should be adapted at the discretion of the veterinarian according to the specific conditions of each animal, region and/or environment, and the recommendations of the local health authority.

DIRECTIONS FOR USE AND ADMINISTRATION:

Vaccination is recommended for animals in good general health, with proper nutritional development, that have been dewormed, are free from fever, and are not experiencing stress due to adverse environmental conditions.

Shake the vial before use. Using a sterile syringe and needle, aseptically withdraw the contents and administer 1 mL subcutaneously.

PRECAUTIONS- WARNINGS- CONTRAINDICATIONS:

Store the product in its original container at 2°C to 8°C. Do not freeze. Use the entire contents immediately after reconstitution. Use sterile disposable needles and syringes for administration. Avoid using disinfectants to clean needles, as they may inactivate the vaccine. Avoid vaccinating pregnant females.
Controlled laboratory tests have demonstrated that this product meets the safety and efficacy standards for immunizing healthy animals. Its effectiveness may be affected by individual animal conditions, which should be taken into consideration by the veterinarian prior to immunization.

ADVERSE REACTIONS

- All vaccines may cause hypersensitivity reactions. In the event of an anaphylactic or hypersensitivity reaction following vaccination, the veterinarian should administer the appropriate treatment.

WITHDRAWAL PERIOD:

- Not applicable.

COMMERCIAL PRESENTATIONS:

- Box of 25 doses.
- Box of 50 doses.